Author Index

Abad, C., see Pérez-Payá, E. 229
Arvidsson, E.
——, Crommen, J., Schill, G. and Westerlund, D. Indirect detection in reversed-phase liquid chromatography. Response deviations in ion-pairing systems 429
Aussems, M., see Duchateau, A. 419
Baba, Y., see Kiniwa, H. 397
Bao, Z., see Yang, S. K. 377
Basselier, J. J., see Desbene, P. L. 305
Bassler, B. J.
——, Kaliszan, R. and Hartwick, R. A. Retention mechanisms on metallic stationary phases 139
Biggs, W. R., see Jinno, K. 209
Blo, G., see Dondi, F. 281
Bongers, J., see Duchateau, A. 419
Braco, L., see Pérez-Payá, E. 229
Brayan, J. G., see Haddad, P. R. 337
Brent, D. A., see Minick, D. J. 177
Bush, H., see Derrico, E. M. 35
Campbell, E. L., see Kroeff, E. P. 45
Campos, A., see Pérez-Payá, E. 229
Carlshaf, A.
—— and Jönsson, J. Å. Gradient elution in hollow-fibre flow field-flow fractionation 89
Cecconie, T., see Thormann, W. 95
Chen, Y.-F., see Meng, Q. C. 271
Cohen, J. R., see Stein, T. A. 267
Cox, G. B., see Gentilucci, T. J. 63
Crombach, M., see Duchateau, A. 419
Crommen, J., see Arvidsson, E. 429
David, P. A.
—— and Novotny, M. Analysis of steroids by capillary supercritical fluid chromatography with phosphorus-selective detection 111
Delucas, L. J., see Meng, Q. C. 271
Derrico, E. M.
——, Bush, H. and Finnegan, P. M. Dioxanyl hydroperoxide in preparative liquid chromatography 35
Desbene, P. L.
——, Desmazieres, B., Basselier, J. J. and Desbene-Monvernay, A. Analytical study of non-ionic surfactants used in enhanced oil recovery. Optimization of analytical conditions in reversed-phase partition chromatography 305
Desbene-Monvernay, A., see Desbene, P. L. 305
Desmarchelier, J. M., see Haddad, P. R. 337
Desmazieres, B., see Desbene, P. L. 305
Dhanesar, S. C.
—— and Gisch, D. J. Mobile phase optimization of a urea-linked chiral stationary phase for the high-performance liquid chromatographic separation of optical isomers 407
Dietz, M. L., see Thormann, W. 95
Dilli, S., see Haddad, P. R. 337
Dobashi, A.
——, Dobashi, Y., Ono, T., Hara, S., Saito, M., Higashidate, S. and Yamauchi, Y. Enantiomer resolution of D- and L-α-amino acid derivatives by supercritical fluid chromatography on novel chiral diamide phases with carbon dioxide 121
Dobashi, Y., see Dobashi, A. 121
Dondi, F.
——, Kahie, Y. D., Lodi, G., Blo, G., Pietrogrande, C. and Reschiglian, P. Solvent selectivity effects in reversed-phase high-performance liquid chromatography of flavonoid compounds 281
Dorsey, J. G., see Sentell, K. B. 193
Doster, D. E.
—— and Zentner, M. Separation of thiodiglycol polyethers and related compounds by reversed-phase high-performance liquid chromatography 293
Duchateau, A.
——, Crombach, M., Aussems, M. and Bongers, J. Determination of the enantiomers of α-amino acids and α-amino acid amines by high-performance liquid chromatography with a chiral mobile phase 419
Emmelin, M., see Félix, G. 347
Faure, A., see Félix, G. 347
Félix, G.
——, Thienpont, A., Emmelin, M. and Faure, A. Hydrocarbon group type analysis of petroleum products by high-performance liquid chromatography on organometallic donor/acceptor-bonded silica 347
Fetzer, J. C., see Jinno, K. 209
Finnegan, P. M., see Derrico, E. M. 35
Firestone, M. A., see Thormann, W. 95
Foley, J. P., see Jeansonne, M. S. 149
Fóti, G.
——, Martinez, C. and Kováts, E. sz. Role of unreacted silanols in the adsorption properties of derivatized silica 243
Frenz, J., see Minick, D. J. 177

Fu, P. P., see Lai, J.-S. 327
——, see Von Tugeln, L. S. 315
Gentilucci, T. J.
——, Sivakoff, S. I., Cox, G. B., Stearns, S. D. and Hutchinson, M. W.
Analytical performance of process-scale liquid chromatography columns 63
Gisch, D. J., see Dhanesar, S. C. 407
Golshan-Shirazi, S.
—— and Guiochon, G.
Theoretical study of system peaks and elution profiles for large concentration bands in the case of a binary eluent containing a strongly sorbed additive 1
—— and Guiochon, G.
Experimental study of system peaks and elution profiles for large concentration bands in the case of a binary eluent containing a strongly sorbed additive 19
Griffiths, P. R., see Jinno, K. 209
Guiochon, G., see Golshan-Shirazi, S. 1, 19
Haddad, P. R.
——, Brayan, J. G., Sharp, G. J., Dilli, S. and Desmarchelier, J. M.
Determination of pyrethroid residues on paddy rice by reversed-phase high-performance liquid chromatography 337
Hara, S., see Dobashi, A. 121
Hartwick, R. A., see Bassler, B. J. 139
Higashidate, S., see Dobashi, A. 121
Hung, S. S., see Lai, J.-S. 327
Hutchinson, M. W., see Gentilucci, T. J. 63
Ibuki, T., see Jinno, K. 209
Ishida, T., see Kiniwa, H. 397
Jeansonne, M. S.
—— and Foley, J. P.
Measurement of statistical moments of resolved and overlapping chromatographic peaks 149
Jinno, K.
——, Ibuki, T., Tanaka, N., Okamoto, M., Fetzer, J. C., Biggs, W. R., Griffiths, P. R. and Olinger, J. M.
Retention behaviour of large polycyclic aromatic hydrocarbons in reversed-phase liquid chromatography on a polymeric octadecylsilica stationary phase 209
Jönsson, J. Å., see Carlshaf, A. 89
Johnson, R. D., see Kroeff, E. P. 45
Jung, H., see Lai, J.-S. 327
Kahie, Y. D., see Dondi, F. 281
Kaliszan, R., see Bassler, B. J. 139
Kasai, M., see Noguchi, K. 365
Katayama, B., see Noguchi, K. 365
Katoh, H., see Kiniwa, H. 397
King, S. J., see Meng, Q. C. 271
Kiniwa, H.
——, Baba, Y., Ishida, T. and Katoh, H.
General evaluation and application to trace analysis of a chiral column for ligand-exchange chromatography 397
Kováts, E. sz., see Fóti, G. 243
Kroeff, E. P.
——, Owens, R. A., Campbell, E. L., Johnson, R. D. and Marks, H. I.
Production scale purification of biosynthetic human insulin by reversed-phase high-performance liquid chromatography 45
Lai, J.-S.
——, Hung, S. S., Unruh, L. E., Jung, H. and Fu, P. P.
Separation of amino- and acetylamino-polycyclic aromatic hydrocarbons by reversed- and normal-phase high-performance liquid chromatography 327
Li, F.
——, Lim, C. K. and Peters, T. J.
Preparation, high-performance liquid chromatographic separation and characterization of hexacarboxylic porphyrinogens 353
Lim, C. K.
—— and Peters, T. J.
Isocratic reversed-phase high-performance liquid chromatography of ribonucleotides, deoxynucleotides, cyclic nucleotides and deoxycyclic nucleotides 259
——, see Li, F. 353
Litzén, A., see Wahlund, K.-G. 73
Lodi, G., see Dondi, F. 281
Lu, X.-L., see Yang, S. K. 377
Luetje, H.
High-performance liquid chromatography of oxidation products of 2,6-dimethylphenol 361
Mandell, C., see Stein, T. A. 267
Marks, H. I., see Kroeff, E. P. 45
Martinez, C., see Fóti, G. 243
Martire, D. E.
Generalized treatment of spatial and temporal column parameters, applicable to gas, liquid and supercritical fluid chromatography. I. Theory 165
Meng, Q. C.
——, King, S. J., Chen, Y.-F., Delucas, L. J. and Oparil, S.
Purification angiotensin-converting enzyme by gel high-performance liquid chromatography 271
Minick, D. J.
——, Brent, D. A. and Frenz, J.
Modeling octanol–water partition coefficients by reversed-phase liquid chromatography 177

Mosher, R. A., see Thormann, W. 95
Mushtaq, M., see Yang, S. K. 377
Noguchi, K.
——, Yanagihara, Y., Kasai, M. and Katayama, B.
Chromatographic properties of a vinyl alcohol copolymer gel column for the analysis of non-ionic surfactants 365
Novotny, M., see David, P. A. 111
Okamoto, M., see Jinno, K. 209
Olinger, J. M., see Jinno, K. 209
Ono, T., see Dobashi, A. 121
Oparil, S., see Meng, Q. C. 271
Oppenheimer, L. E.
—— and Smith, G. A.
Sedimentation field-flow fractionation of colloidal metal hydrosols 103
Owens, R. A., see Kroeff, E. P. 45
Pérez-Payá, E.
——, Braco, L., Campos, A., Soria, V. and Abad, C.
Solution properties of polyelectrolytes. IV. Use of a new hydrophilic size-exclusion chromatographic packing for the separation of anionic and cationic polyions 229
Peters, T. J., see Li, F. 353
——, see Lim, C. K. 259
Pietrogrande, C., see Dondi, F. 281
Rennie, P. J.
Determination of phenylalanine in river water by high-performance liquid chromatography 277
Reschiglian, P., see Dondi, F. 281
Sabio, M., see Topiol, S. 129
Saito, M., see Dobashi, A. 121
Schill, G., see Arvidsson, E. 429
Sentell, K. B.
—— and Dorsey, J. G.
Retention mechanisms in reversed-phase chromatography stationary phase bonding density and solute selectivity 193
Sharp, G. J., see Haddad, P. R. 337
Shou, M., see Yang, S. K. 377
Sivakoff, S. I., see Gentilucci, T. J. 63
Smith, G. A., see Oppenheimer, L. E. 103
Soria, V., see Pérez-Payá, E. 229
Stearns, S. D., see Gentilucci, T. J. 63
Stein, T. A.
——, Cohen, J. R., Mandell, C. and Wise, L.
Determination of elastase activity by reversed-phase high-performance liquid chromatography 267
Tanaka, N., see Jinno, K. 209
Thienpont, A., see Félix, G. 347
Thormann, W.
——, Firestone, M. A., Dietz, M. L., Cecconie, T. and Mosher, R. A.
Focusing counterparts of electrical field flow fractionation and capillary zone electrophoresis. Electrical hyperlayer field flow fractionation and capillary isoelectric focusing 95
Topiol, S.
—— and Sabio, M.
Computational chemical studies of chiral stationary phase models. Complexes of methyl N-(2-naphthyl)alaninate with N-(3,5-dinitrobenzoyl)leucine *n*-propylamide 129
Tungeln, L. S., von, see Von Tungeln, L. S. 315
Unruh, L. E., see Lai, J.-S. 327
Von Tungeln, L. S.
—— and Fu, P. P.
High-performance liquid chromatographic separation of ring-oxidized metabolites of nitro-polycyclic aromatic hydrocarbons 315
Wahlund, K.-G.
—— and Litzén, A.
Application of an asymmetrical flow field-flow fractionation channel to the separation and characterization of proteins, plasmids, plasmid fragments, polysaccharides and unicellular algae 73
Weems, H. B., see Yang, S. K. 377
Westerlund, D., see Arvidsson, E. 429
Wise, L., see Stein, T. A. 267
Yamauchi, Y., see Dobashi, A. 121
Yanagihara, Y., see Noguchi, K. 365
Yang, S. K.
——, Mushtaq, M., Bao, Z., Weems, H. B., Shou, M. and Lu, X.-L.
Improved enantiomeric separation of dihydrodiols of polycyclic aromatic hydrocarbons on chiral stationary phases by derivatization to *O*-methyl ethers 377
Zentner, M., see Doster, D. E. 293